



On Rational and Irrational Values of Trigonometric Functions

Veli Akarsu*¹ 

¹Zonguldak Bülent Ecevit University, Zonguldak Vocational School, Department of Architecture and Urban Planning, 67970, Zonguldak, Türkiye; (aveli9827@gmail.com)

Keywords

Trigonometry,
Trigonometric Functions,
Rational Number,
Irrational Number.

Research Article

Received : 20.04.2023
Revised : 08.08.2023
Accepted : 12.09.2023
Published : 30.09.2023

* Corresponding Author
aveli9827@gmail.com



Abstract

The beginning of trigonometry goes back 4000 years from today. Today, there are Euclidean and Non-Euclidean trigonometry. It can be said that trigonometry emerged from the need to make maps containing the position information of the stars, to determine the time and to make a calendar, mostly in astronomy. Each of the six trigonometric functions is defined according to the directed plane angle. For each angle in the domain of these functions, their values correspond to a real number consisting of rational or irrational numbers. Trigonometric functions have an important position in basic sciences and technology as well as calculations in engineering and architecture. In addition to being a branch of mathematics, trigonometry is widely used in solving geometry and analysis problems. It has an indispensable importance in engineering and architectural design and calculations. The values of trigonometric functions are usually an irrational number, with the exception of some special angle values. Irrational numbers are numbers that are not proportional. In other words, they are numbers whose results are uncertain. Examples are numbers such as pi, e, and radical. The irrational values of trigonometric functions, which are infinite decimal expansions, are given in the tables by rounding them to only four or six digits. When entering the trigonometric function values in the tables (or the values obtained in the libraries of electronic calculators) into algebraic operations, the resulting numbers are approximated. In the article, besides showing that most of the trigonometric function values such as $\sin\theta$, $\cos\theta$ and $\tan\theta$ of many θ angles are irrational, the effect of these functions on the result values of calculations made in engineering and architecture is interpreted.

1. Introduction

In ancient Babylon, geometry arose from the practical needs of rulers, cartographers, and builders (Britton et al. 2011). From measurements of fields, walls, posts, buildings, gardens, canals, and ziggurats (temple tower of the city god in Mesopotamia), squares, rectangles, trapezoids, and right triangles were used as the main types of practical geometric shapes (Mansfield and Wildberger, 2017). Ratio-based measurements were an important measurement in Egyptian architecture used to describe the reverse slope ($\cot\theta$) of the pyramids.

Evidence that the foundations of trigonometry date back to the ancient Babylonian period between the 19th and 16th centuries BC was reached by deciphering the Plimpton 322 (P322) tablet, the most advanced scientific

work of antiquity. The P322 tablet is a trigonometric table thousands of years ahead of its time (Mansfield and Wildberger, 2017).

The table of values of modern trigonometric functions consists of rational and / or irrational ratios of trigonometric functions $\sin\theta$, $\cos\theta$ and $\tan\theta$, which show the ratio of the side lengths of right triangles with hypotenuse side length, 1 unit.

The only correct trigonometric table of the world is the trigonometric table of the astronomer and mathematician Hipparchus (190-120 BC) (Mansfield and Wildberger, 2017). The pioneer of studies on trigonometry was Hipparchus, and her successor was Ptolemy (85-160 AD) (Stewart, 2009). The first person whose trigonometric table has survived is Alexandrian

Cite this;

Akarsu, V. (2023). On Rational and Irrational Values of Trigonometric Functions. *Advanced Geomatics*, 3(2), 56-62.

astronomer Claudius Ptolemaios (Ptolemy), who lived in the 2nd century AD (Van Brummelen, 2012). Ptolemy wrote a 13-volume book on astronomy called *Mathēmatikē Syntaxis*. This book has been translated into Arabic as the *Almagest*. In the first volume of the book, the relationship between trigonometry and astronomy was examined, and he arranged the chords ruler with the method he developed for calculating the chords of angles (Gölgeleyen ve Akarsu, 2022). He calculated the approximate value of the beam corresponding to an angle of one degree. The table of values of the sine trigonometric function is included in the book.

It can be said that the need for table of values of trigonometric functions is in astronomical observations and calendar making works rather than solving geometry problems. Each of the six trigonometric functions, consisting of three basic ($\sin\theta$, $\cos\theta$, $\tan\theta$) and three auxiliary ($\csc\theta$, $\sec\theta$, $\cot\theta$) trigonometric functions, is defined according to the directed plane angle. For each angle in the domain of these functions, their values in the value set correspond to a real number consisting of rational and/or irrational numbers. The first discovered irrational number is $\sqrt{2}$. This discovery was made because of the curiosity of Greek geometers to measure the diagonal length of a square with a side length of 1 unit with a rational number. With the Pythagorean theorem, it has been proved that there is no rational number satisfying $x^2=1+1=2$. In the decimal notation of $\sqrt{2}$, it is followed by an infinite number of digits after the decimal symbol. Likewise, today, π has been calculated to tens of trillions of digits after the decimal point. However, the basic digits of the decimal number system used today are based on the Dedekind-Peano axioms system for natural numbers. By expanding the natural numbers axioms system, real and complex number systems are obtained. Although the decimal representation of every rational number consists of a limited number of digits, this expansion continues indefinitely for irrational numbers.

In the article, besides showing that most of the basic trigonometric function values such as $\sin\theta$, $\cos\theta$ and $\tan\theta$ of many θ plane angles are irrational, the effect of these values on the result values of calculations made in science is examined.

2. Method

2.1. Some trigonometric identities and equations

For plane angles α and β ,

$$\begin{aligned} \cos(\alpha + \beta) &= \cos\alpha \cdot \cos\beta - \sin\alpha \cdot \sin\beta \\ \sin(\alpha + \beta) &= \sin\alpha \cdot \cos\beta + \cos\alpha \cdot \sin\beta \end{aligned} \quad (1)$$

$$\begin{aligned} \cos(\alpha - \beta) &= \cos\alpha \cdot \cos\beta + \sin\alpha \cdot \sin\beta \\ \sin(\alpha - \beta) &= \sin\alpha \cdot \cos\beta - \cos\alpha \cdot \sin\beta \end{aligned} \quad (2)$$

Matrices enable us to write them in a compact form,

$$\begin{bmatrix} \cos\alpha & \sin\alpha \\ -\sin\alpha & \cos\alpha \end{bmatrix} \begin{bmatrix} \cos\beta & \sin\beta \\ -\sin\beta & \cos\beta \end{bmatrix} = \begin{bmatrix} \cos(\alpha + \beta) & \sin(\alpha + \beta) \\ -\sin(\alpha + \beta) & \cos(\alpha + \beta) \end{bmatrix} \quad (1a)$$

$$\begin{bmatrix} \cos\alpha & \sin\alpha \\ \sin\alpha & -\cos\alpha \end{bmatrix} \begin{bmatrix} \cos\beta & \sin\beta \\ \sin\beta & -\cos\beta \end{bmatrix} = \begin{bmatrix} \cos(\alpha - \beta) & \sin(\alpha - \beta) \\ \sin(\alpha - \beta) & \cos(\alpha - \beta) \end{bmatrix} \quad (2a)$$

equations can be written (Beasley, 1858; Dörrie, 1950; Daut, 1951; Helton, 1972; Durell, 1975; Sigl, 1977; Barnet, 1991; Akarsu 2005; Akarsu, 2009).

Also, for any plane angle α ,

$$\sin^2\alpha + \cos^2\alpha = 1 \quad (3)$$

the trigonometric equivalent of the Pythagorean theorem can be written.

If $\alpha = \beta = \theta$ is taken and substituted in equations (1) and (2) and equation (3) is taken into account,

$$\begin{aligned} \cos 2\theta &= \cos^2\theta - \sin^2\theta = 2\cos^2\theta - 1 \\ &= 1 - 2\sin^2\theta \end{aligned} \quad (4)$$

$$\sin 2\theta = 2\sin\theta \cdot \cos\theta \quad (5)$$

equations (4) and (5) can be obtained.

Similarly, if $\beta = 2\theta$ is written in equation (1) and equations (3), (4) and (5) are taken into account,

$$\cos 3\theta = 4 \cdot \cos^3\theta - 3\cos\theta \quad (6)$$

equation (6) can also be written (Beasley, 1858; Dörrie, 1950; Daut, 1951; Durell, 1975).

$$\sin 2\theta = 2\sin\theta \cdot \cos\theta$$

$$\sin 3\theta = 3 \cdot \sin\theta \cdot \cos^2\theta - \sin^3\theta$$

$$\sin 4\theta = 4 \cdot \sin\theta \cdot \cos^3\theta - 4\cos\theta \cdot \sin^3\theta \quad (6a)$$

$$\begin{aligned} \sin 5\theta &= 5 \cdot \sin\theta - 20 \cdot \sin^3\theta + 16 \cdot \sin^5\theta \\ &\vdots \end{aligned}$$

Or

$$\sin 2\theta = 2\sin\theta \cdot \cos\theta$$

$$\sin 3\theta = 2\cos\theta \cdot \sin 2\theta - \sin\theta$$

$$\sin 4\theta = 2\cos\theta \cdot \sin 3\theta - \sin 2\theta \quad (6b)$$

$$\begin{aligned} \sin 5\theta &= 2\cos\theta \cdot \sin 4\theta - \sin 3\theta \\ &\vdots \end{aligned}$$

$$\begin{aligned} \cos 2\theta &= \cos^2\theta - \sin^2\theta = 2 \cdot \cos^2\theta - 1 \\ &= 1 - 2 \cdot \sin^2\theta \end{aligned}$$

$$\cos 3\theta = \cos^3\theta - 3 \cdot \cos\theta \cdot \sin^2\theta$$

$$\cos 4\theta = \cos^4\theta - 6 \cdot \cos^2\theta \cdot \sin^2\theta + \sin^4\theta \quad (6c)$$

$$\begin{aligned} \cos 5\theta &= 5 \cdot \cos\theta - 20 \cdot \cos^3\theta + 16 \cdot \cos^5\theta \\ &\vdots \end{aligned}$$

or

$$\begin{aligned}
 \cos 2\theta &= 2 \cdot \cos^2 \theta - 1 \\
 \cos 3\theta &= 2 \cos \theta \cdot \cos 2\theta - \cos \theta \\
 \cos 4\theta &= 2 \cos \theta \cdot \cos 3\theta - \cos 2\theta \\
 \cos 5\theta &= 2 \cos \theta \cdot \cos 4\theta - \cos 3\theta \\
 &\vdots \\
 \tan 2\theta &= \frac{2 \cdot \tan \theta}{1 - \tan^2 \theta} \\
 \tan 3\theta &= \frac{3 \cdot \tan \theta - \tan^3 \theta}{1 - 3 \cdot \tan^2 \theta} \\
 \tan 4\theta &= \frac{4 \cdot \tan \theta - 4 \tan^3 \theta}{1 - 6 \cdot \tan^2 \theta + \tan^4 \theta} \\
 &\vdots
 \end{aligned}
 \tag{6d}$$

or

$$\begin{aligned}
 \tan 2\theta &= \frac{2 \cdot \tan \theta}{1 - \tan^2 \theta} \\
 \tan 3\theta &= \frac{\tan \theta + \tan 2\theta}{1 - \tan \theta \cdot \tan 2\theta} \\
 \tan 4\theta &= \frac{\tan \theta + \tan 3\theta}{1 - \tan \theta \cdot \tan 3\theta} \\
 &\vdots
 \end{aligned}
 \tag{6f}$$

From (6a) to (6f), different trigonometric equations can be written (Sigl, 1977).

2.2. Irrationality of Trigonometric Ratios

In this section, the irrationality of the values of the basic trigonometric functions $\cos 20^\circ$, $\sin 10^\circ$ ve $\tan 15^\circ$ of rational angles such as 20° , 10° ve 15° are shown.

2.2.1. Irrationality of $\cos 20^\circ$ Trigonometric Function Value

For $\theta = 20^\circ$ let $\cos 60^\circ = \frac{1}{2}$ and $\cos 20^\circ = x$. If these values are substituted in (6),

$$\begin{aligned}
 \cos 60^\circ &= 4 \cdot \cos^3 20^\circ - 3 \cdot \cos 20^\circ \\
 8x^3 - 6x - 1 &= 0
 \end{aligned}
 \tag{7}$$

(7) set up the polynomial equation. Because of the way the polynomial equation (7) is set up, $\cos 20^\circ$ is a root of the equation. The possible rational roots of equation (7) are $\pm 1, \pm 1/2, \pm 1/4, \pm 1/8$. If these values are substituted in equation (7), it is seen that no root satisfies equation (7).

Since none of the above 8 roots satisfy equation (7), they are not real roots. Therefore, it is understood that equation (7) has no real roots, so $\cos 20^\circ$ is an

irrational number (Jung, 1962; Helton, 1972). Whether $\pm 1, \pm 1/(2, \pm 1/4, \pm 1/8)$ rational roots are the real roots of the equation (7) can also be understood by comparing the value of $\cos 20^\circ = 0,939693$

Namely,

$$\begin{aligned}
 \cos 30^\circ &< \cos 20^\circ < \cos 0^\circ \text{ or} \\
 0,866025 &< \cos 20^\circ < 1
 \end{aligned}
 \tag{8}$$

(8) inequality can be written.

2.2.2. $\sin 10^\circ$ Displaying the Irrationality of Trigonometric

Function Value

In equation (2), $\beta = 2\theta$ is written and equations (3), (4) and (5) are taken into account,

$$\sin 3\theta = 3 \cdot \sin \theta - 4 \cdot \sin^3 \theta
 \tag{9}$$

equation (9) can be written (Niven, 1956; Jung, 1962; Niven, 1964; Helton, 1972; Bergen, 2009; Paolillo and Vincenzi, 2021).

For $\theta = 10^\circ$, let $\sin 30^\circ = \frac{1}{2}$ and $\sin 10^\circ = x$. If these values are substituted in (9),

$$\sin 30^\circ = 3 \cdot \sin 10^\circ - 4 \cdot \sin^3 10^\circ
 \tag{10}$$

$$8x^3 - 6x + 1 = 0$$

(10) polynomial equation can be established.

(4) by virtue of equality,

$$8x^3 - 6x + 1 = 0
 \tag{11}$$

(11) equation is taken into account.

Let $\sin 10^\circ$ be rational in equation (11). In this case, the expressions $\sin^2 10^\circ$ and $1 - 2\sin^2 10^\circ$ would also be rational. Whereas, $\cos 20^\circ$ was shown to be irrational by equation (7) and inequality (8). Therefore, it is concluded that $\sin 10^\circ$ is also irrational in order not to cause a contradiction.

2.2.3. Demonstrating the Irrationality of the $\tan 15^\circ$ Trigonometric Function Value

Starting from equation (4),

$$\tan \theta = \sqrt{\frac{1 - \cos 2\theta}{1 + \cos 2\theta}}
 \tag{12}$$

(12) can be written.

$$\theta = 15^\circ \text{ için } \cos 30^\circ = \frac{\sqrt{3}}{2} \text{ dir. For } \theta = 15^\circ, \cos 30^\circ = \frac{\sqrt{3}}{2}.$$

If these values are substituted in the equation (12);

$$\tan 15^\circ = \sqrt{\frac{2-\sqrt{3}}{2+\sqrt{3}}} = 0,2679491924.$$

Since the value of $\tan 15^\circ = \sqrt{\frac{2-\sqrt{3}}{2+\sqrt{3}}}$ in the square root on the right is irrational, its square root is also irrational. Hence, it is understood that the value of $\tan 15^\circ$ is also irrational. Hence, it is understood that the value of $\tan 15^\circ$ is also irrational.

3. Extending the Irrationality of Trigonometric Function Values

It can also be extended outside of some special angles, such as the indicated irrationality of the values of the trigonometric functions $\cos 20^\circ$, $\sin 10^\circ$ and $\tan 15^\circ$. For example, trigonometric function values of angles like $62^\circ 05' 10''$ are irrational. Integer angles such as 0° , 30° , 45° and 60° and when these angles are added by 90° and its multiples, or subtracted from 90° and its multiples, all trigonometric angles The values of the functions cannot be said to be rational. But at least one trigonometric function value is rational.

For example, for all trigonometric function values of 30° ,

$$\sin 30^\circ = \frac{1}{2}, \csc 30^\circ = 2, \text{ rational; } \cos 30^\circ = \frac{\sqrt{3}}{2},$$

$$\sec 30^\circ = \frac{2}{\sqrt{3}} \text{ irrational; } \tan 30^\circ = \frac{\sqrt{3}}{3}, \cot 30^\circ = \sqrt{3}$$

irrational, It is seen that 2 trigonometric function values are rational and 4 trigonometric function values are irrational.

Starting from equation (4),

$$\begin{aligned} \cos \theta &= \sqrt{\frac{1 + \cos 2\theta}{2}}, \sin \theta = \sqrt{\frac{1 - \cos 2\theta}{2}}, \\ \tan \theta &= \sqrt{\frac{1 - \cos 2\theta}{1 + \cos 2\theta}} \end{aligned} \quad (13)$$

(13) equations can be written. If angle θ is any angle that makes the trigonometric function value $\cos 2\theta$ irrational, the function values $\sin \theta = f(\cos 2\theta)$, $\cos \theta = f(\cos 2\theta)$ can also be defined using the irrationality of $\cos 2\theta$ to define an infinite number of angles θ_i , $-360^\circ \leq \theta_i \leq 360^\circ$ the proof that the values of the trigonometric functions in the range of values $-1 \leq \sin \theta$, $\cos \theta \leq 1$ and $-\infty \leq \tan \theta \leq \infty$, are irrational can be extended. Also, this expansion can be done using the trigonometric equations (6a) to (6f).

$$\sin \frac{\theta}{2} = \pm \sqrt{\frac{1 - \cos \theta}{2}}, \cos \frac{\theta}{2} = \pm \sqrt{\frac{1 + \cos \theta}{2}}$$

The $\theta = 90^\circ$ angle can be taken as half of the continuous and the irrational values of the trigonometric functions can be calculated.

$$\sin 45^\circ = \frac{\sqrt{2}}{2}, \cos 45^\circ = \frac{\sqrt{2}}{2}$$

$$\sin 22,5^\circ = \frac{1}{2}\sqrt{2 - \sqrt{2}}, \cos 22,5^\circ = \frac{1}{2}\sqrt{2 + \sqrt{2}}$$

$$\sin 11,25^\circ = \frac{1}{2}\sqrt{2 - \sqrt{2 + \sqrt{2}}}$$

$$\cos 11,25^\circ = \frac{1}{2}\sqrt{2 + \sqrt{2 + \sqrt{2}}}$$

$$\sin 5,625^\circ = \frac{1}{2}\sqrt{2 - \sqrt{2 + \sqrt{2 + \sqrt{2}}}}$$

$$\cos 5,625^\circ = \frac{1}{2}\sqrt{2 + \sqrt{2 + \sqrt{2 + \sqrt{2}}}}$$

irrational values can be calculated.

Similarly, taking $\theta = \frac{90^\circ}{2^n}$, $n \in \mathbb{Z}^+$, the values of the trigonometric functions of the angles are,

$$\sin \frac{90^\circ}{2^n} = \frac{1}{2}\sqrt{2 - \sqrt{2 + \sqrt{2 + \sqrt{2} + \dots}}}$$

$$\cos \frac{90^\circ}{2^n} = \frac{1}{2}\sqrt{2 + \sqrt{2 + \sqrt{2 + \sqrt{2} + \dots}}}$$

it can be calculated by chaining equations (Sigl, 1977).

$\cos \theta = t$ olmak üzere,

$$\sin \frac{\theta}{2} = \sqrt{\frac{1-t}{2}} = \frac{1}{2}\sqrt{2-2t}, \cos \frac{\theta}{2} = \sqrt{\frac{1+t}{2}} = \frac{1}{2}\sqrt{2+2t}$$

$$\cos \frac{\theta}{2} = \frac{1}{2}\sqrt{2+2t} = k \text{ including,}$$

$$\sin \frac{\theta}{4} = \frac{1}{2}\sqrt{2-2k} = \frac{1}{2}\sqrt{2-\sqrt{2+2t}}$$

$$\cos \frac{\theta}{4} = \frac{1}{2}\sqrt{2+2k} = \frac{1}{2}\sqrt{2+\sqrt{2+2t}}$$

equations can be written.

$$\begin{aligned}\cos \frac{\theta}{4} &= \frac{1}{2} \sqrt{2 + \sqrt{2 + 2k}} = r \\ \sin \frac{\theta}{8} &= \frac{1}{2} \sqrt{2 - 2r} = \frac{1}{2} \sqrt{2 - \sqrt{2 + \sqrt{2 + 2t}}} \\ \cos \frac{\theta}{8} &= \frac{1}{2} \sqrt{2 + 2r} = \frac{1}{2} \sqrt{2 + \sqrt{2 + \sqrt{2 + 2t}}}\end{aligned}\quad (14)$$

$\cos \theta = t$ including,

$$\begin{aligned}\sin \frac{\theta}{2} &= \frac{1}{2} \sqrt{2 - 2t} \\ \sin \frac{\theta}{4} &= \frac{1}{2} \sqrt{2 - \sqrt{2 + 2t}} \\ \sin \frac{\theta}{8} &= \frac{1}{2} \sqrt{2 - \sqrt{2 + \sqrt{2 + 2t}}} \\ \cos \frac{\theta}{2} &= \frac{1}{2} \sqrt{2 + 2t} \\ \cos \frac{\theta}{4} &= \frac{1}{2} \sqrt{2 + \sqrt{2 + 2t}} \\ \cos \frac{\theta}{8} &= \frac{1}{2} \sqrt{2 + \sqrt{2 + \sqrt{2 + 2t}}}\end{aligned}\quad (15)$$

equations (14) and (15) can be written.

Now, using the equations (14), (15) (1) and (2), the irrationality of the sine and cosine trigonometric function values of many angles is shown below.

$$\sin 30^\circ = \frac{1}{2}, \cos 30^\circ = \frac{1}{2} \sqrt{3}$$

$$\sin 15^\circ = \frac{1}{2} \sqrt{2 - \sqrt{3}}$$

$$\cos 15^\circ = \frac{1}{2} \sqrt{2 + \sqrt{3}}$$

$$\sin 18^\circ = \frac{1}{4} (\sqrt{5} - 1)$$

$$\cos 18^\circ = \frac{1}{4} \sqrt{10 + 2\sqrt{5}}$$

$$\sin 7,5^\circ = \frac{1}{2} \sqrt{2 - \sqrt{2 + \sqrt{3}}}$$

$$\cos 7,5^\circ = \frac{1}{2} \sqrt{2 + \sqrt{2 + \sqrt{3}}}$$

$$\sin 3,75^\circ = \frac{1}{2} \sqrt{2 - \sqrt{2 + \sqrt{2 + \sqrt{3}}}}$$

$$\cos 3,75^\circ = \frac{1}{2} \sqrt{2 + \sqrt{2 + \sqrt{2 + \sqrt{3}}}}$$

$$\begin{aligned}\sin 3^\circ &= \sin(18^\circ - 15^\circ) \\ &= \sin 18^\circ \cos 15^\circ - \cos 18^\circ \sin 15^\circ \\ &= \frac{1}{16} (\sqrt{30} + \sqrt{10} - \sqrt{6} - \sqrt{2} \\ &\quad - 2\sqrt{20 + 4\sqrt{5} - 10\sqrt{3} - 2\sqrt{15}})\end{aligned}$$

$$\begin{aligned}\sin 6^\circ &= \cos 84^\circ = \sin(36^\circ - 30^\circ) \\ &= \frac{1}{8} (\sqrt{30 - 6\sqrt{5} - \sqrt{5} - 1})\end{aligned}$$

$$\begin{aligned}\sin 9^\circ &= \cos 81^\circ = \sin(45^\circ - 36^\circ) \\ &= \frac{1}{8} (\sqrt{10} + \sqrt{2} - 2\sqrt{5 - \sqrt{5}})\end{aligned}$$

$$\begin{aligned}\sin 12^\circ &= \cos 78^\circ = \sin(30^\circ - 18^\circ) \\ &= \frac{1}{8} (\sqrt{10 + 2\sqrt{5} - \sqrt{15} + \sqrt{3}})\end{aligned}$$

$$\sin 21^\circ = \cos 69^\circ = \sin(36^\circ - 15^\circ)$$

$$\begin{aligned}&= \frac{1}{8} (\sqrt{15 - 3\sqrt{5}} + \sqrt{5 - \sqrt{5}} - \sqrt{10 - 5\sqrt{3}} \\ &\quad - \sqrt{2 - \sqrt{3}})\end{aligned}$$

$$\sin 24^\circ = \cos 66^\circ = \sin(54^\circ - 30^\circ)$$

$$= \frac{1}{8} (\sqrt{15} + \sqrt{3} - \sqrt{10 - 2\sqrt{5}})$$

$$\sin 27^\circ = \cos 63^\circ = \sin(45^\circ - 18^\circ)$$

$$= \frac{1}{8} (2\sqrt{5 + \sqrt{5}} - \sqrt{10} + \sqrt{2})$$

$$\begin{aligned}\sin 33^\circ &= \cos 57^\circ \\ &= \sin(15^\circ + 18^\circ) \\ &= \frac{1}{16} (2\sqrt{20 + 4\sqrt{5} - 10\sqrt{3} - 2\sqrt{15}} + \sqrt{30} + \sqrt{10} \\ &\quad - \sqrt{6} - \sqrt{2})\end{aligned}$$

$$\begin{aligned}\sin 39^\circ &= \cos 51^\circ = \sin(54^\circ - 15^\circ) \\ &= \frac{1}{16}(\sqrt{30} + \sqrt{6} + \sqrt{10} + \sqrt{2} \\ &\quad - 2\sqrt{20 + 2\sqrt{15} - 4\sqrt{5} - 10\sqrt{3}})\end{aligned}$$

$$\begin{aligned}\sin 42^\circ &= \cos 48^\circ = \sin(60^\circ - 18^\circ) \\ &= \frac{1}{8}(\sqrt{30 + 6\sqrt{5}} - \sqrt{5} + 1)\end{aligned}$$

$$\begin{aligned}\sin 48^\circ &= \cos 42^\circ = \sin(30^\circ + 18^\circ) \\ &= \frac{1}{8}(\sqrt{30 + 2\sqrt{5}} + \sqrt{15} - \sqrt{3})\end{aligned}$$

$$\begin{aligned}\sin 51^\circ &= \cos 39^\circ = \sin(36^\circ + 15^\circ) \\ &= \frac{1}{8}(\sqrt{15 - 3\sqrt{5}} - \sqrt{5 - \sqrt{5}} \\ &\quad + \sqrt{10 - 5\sqrt{3}} + \sqrt{2 - \sqrt{3}})\end{aligned}$$

$$\begin{aligned}\sin 57^\circ &= \cos 33^\circ = \sin(72^\circ - 15^\circ) \\ &= \frac{1}{8}(\sqrt{15 + 3\sqrt{5}} - \sqrt{5 + \sqrt{5}} \\ &\quad - \sqrt{10 - 5\sqrt{3}} + \sqrt{2 - \sqrt{3}})\end{aligned}$$

$$\begin{aligned}\sin 63^\circ &= \cos 27^\circ = \sin(45^\circ + 18^\circ) \\ &= \frac{1}{8}(2\sqrt{5 + \sqrt{5}} + \sqrt{10} - \sqrt{2})\end{aligned}$$

$$\begin{aligned}\sin 66^\circ &= \cos 24^\circ = \sin(36^\circ + 30^\circ) \\ &= \frac{1}{8}(\sqrt{30 - 6\sqrt{5}} + \sqrt{5} + 1)\end{aligned}$$

$$\begin{aligned}\sin 69^\circ &= \cos 21^\circ = \sin(54^\circ + 15^\circ) \\ &= \frac{1}{16}(\sqrt{30} + \sqrt{10} + \sqrt{6} + \sqrt{2} \\ &\quad + 2\sqrt{20 + 2\sqrt{15} - 4\sqrt{5} - 10\sqrt{3}})\end{aligned}$$

$$\begin{aligned}\sin 78^\circ &= \cos 12^\circ = \sin(60^\circ + 18^\circ) \\ &= \frac{1}{8}(\sqrt{30 + 6\sqrt{5}} + \sqrt{5} - 1)\end{aligned}$$

$$\begin{aligned}\sin 81^\circ &= \cos 9^\circ = \sin(45^\circ + 36^\circ) \\ &= \frac{1}{8}(\sqrt{10} + \sqrt{2} + 2\sqrt{5 - \sqrt{5}})\end{aligned}$$

$$\begin{aligned}\sin 84^\circ &= \cos 6^\circ = \sin(54^\circ + 30^\circ) \\ &= \frac{1}{8}(\sqrt{10 - 2\sqrt{5}} + \sqrt{15} + \sqrt{3})\end{aligned}$$

$$\begin{aligned}\sin 87^\circ &= \cos 3^\circ = \sin(72^\circ + 15^\circ) \\ &= \frac{1}{8}(\sqrt{15 + 3\sqrt{5}} + \sqrt{5 + \sqrt{5}} \\ &\quad + \sqrt{10 - 5\sqrt{3}} - \sqrt{2 - \sqrt{3}})\end{aligned}$$

As shown in the above equations, $3^\circ, 6^\circ, 9^\circ, 12^\circ, 15^\circ, 18^\circ, 21^\circ, 24^\circ, 27^\circ, 33^\circ, 39^\circ, 42^\circ, 48^\circ, 51^\circ, 57^\circ, 63^\circ, 66^\circ, 69^\circ, 78^\circ, 81^\circ, 84^\circ, 87^\circ$ sine and cosine trigonometric function values of angles, $\sqrt{2}, \sqrt{3}, \sqrt{5}, \sqrt{10}, \sqrt{15}, \sqrt{30}$ as they are calculated based on irrational numbers, their results are irrational (Beasley, 1858; Dörrie, 1950; Daut, 1951; Niven, 1956; Jung, 1962; Niven, 1964; Helton, 1972; Durell, 1975; Sigl, 1977; Barnett, 1991; Akarsu 2005; Akarsu, 2009; Bergen, 2009; Paolillo and Vincenzi, 2021). The rational and irrational values of trigonometric functions are used in applications of engineering sciences (Maune, 2001; Yakar, 2011; Pellicani et al. 2016; Alptekin et al. 2019; Ulvi et al., 2019; Alptekin, 2020).

4. Results and Discussion

Trigonometric functions are periodic functions with important applications in mathematics, science and technology. Trigonometric functions are widely used in science. Although rational numbers are numbers with finite decimal expansions, irrational numbers have infinite or noncyclic expansions. It has been shown by equations (13) that the trigonometric functions have infinitely many values in the $0^\circ \leq \alpha \leq 360^\circ$ definition range, while they have infinitely many irrational values in the $-1 \leq \sin \alpha, \cos \alpha \leq 1$ and $-\infty \leq \tan \alpha \leq \infty$ value ranges. As it can be understood from the examples given, it is understood from the examples (2.2.1), (2.2.2) and (2.2.3) above that although the rational values of the trigonometric functions in the value set are quite few, the majority of them consist of irrational numbers. Intermediate calculation values using trigonometric functions in basic sciences, astronomy, engineering and architecture, and technology contain rounding errors. Therefore, it is thought that it is important to be aware that the results of the calculations made with the irrational values of the trigonometric functions used in science always contain an error. It can be said that trigonometric functions, which are a simple geometric way of thinking, contribute to the development of today's science, mathematics, engineering and architectural education.

Author contributions

The article is single-authored.

Conflicts of interest

The authors declare no conflicts of interest.

Statement of Research and Publication Ethics

Research and publication ethics were complied with in the study.

References

- Akarsu, V. (2005). Geometride, Uzay, Düşey ve Yatay Açılar Arasındaki Fonksiyonel İlişki, *Selcuk University Journal of Engineering Sciences*, 4(3), 134-142.
- Akarsu, V. (2009). Düzlem Üçgende Açıların Kenarlardan Bulunması, *Erciyes Üniversitesi Fen Bilimleri Enstitüsü Fen Bilimleri Dergisi*, 25(1), 390-399.
- Alptekin, A., & Yakar, M (2020). Heyelan bölgesinin İHA kullanarak modellenmesi. *Türkiye İnsansız Hava Araçları Dergisi*, 2(1), 17-21.
- Alptekin, A., Çelik, M. Ö., Doğan, Y., & Yakar, M. (2019). Mapping of a rockfall site with an unmanned aerial vehicle. *Mersin Photogrammetry Journal*, 1(1), 12-16.
- Barnet, R. A. (1991). *Analytic Trigonometry with Applications*, Wadsworth Publishing Company, Belmont, California, 431.
- Beasley, R. D. (1858). *Plane Trigonometry*, Cambridge: Macmillan and Co., London, 106.
- Bergen, J. (2009). Values of Trigonometric Functions, *Math Horizons*, 16(3), 17-19, DOI: 10.1080/10724117.2009.11974811
- Britton, J. P., Proust, C., & Shnider, S. (2011). Plimpton 322: a review and a different perspective. *Archive for history of exact sciences*, 65, 519-566.
- Daut, W. (1951). *Ebene Trigonometrie*, Paedagogischer Verlag Berthold Schulz, Berlin-Hannover-Frankfurt/Mein, 212.
- Dörrie, H. (1950). *Ebene und Sphaerische Trigonometrie*, Verlag Von R. Oldenbourg, München, 518s.
- Durell, C. V. (1975). *Matriculation Trigonometry*, G. Bell and Sons, Ltd, London, 151s.
- Gölgeleyen F., & Akarsu V (2022). İnsan Aklının Evrimi: Lebombo Mağarası'ndan Mouseion'a Matematiğin Serüveni, Türkiye'de Mühendislik ve Mimarlığın 250 Yılı Uluslararası Sempozyumu, 19-21 Mayıs, İTÜ, İstanbul.
- Helton, F. F. (1972). *Analytic Trigonometry*, W.B. Saunders Company, Philadelphia-London-Toronto, 297.
- Jung, H. W. E. (1962). Sayılar Teorisine Giriş, *Türk Matematik Derneği Yayınları*, Çeviren: Orhan Ş. İçen, 176.
- Mansfield, D. F., & Wildberger, N. J. (2017). Plimpton 322 is Babylonian exact sexagesimal trigonometry. *Historia Mathematica*, 44(4), 395-419.
- Maune, D. F. (2001) *Digital elevation model technologies and applications: The DEM User manual*. The American Society for Photogrammetry and Remote Sensing. ISBN:1-57083-064-9
- Niven, I. M. (1956). *Irrational Numbers*. Wiley, New York.
- Niven, I. M. (1964). *Rasyonel ve İrrasyonel Sayılar*, Çeviri: Adnan Kırıl, *Türk matematik Derneği Yayınları*, 192s.
- Paolillo, B., & Vincenzi, G. (2021). On the rational values of trigonometric functions of angles that are rational in degrees. *Mathematics Magazine*, 94(2), 132-134.
- Pellicani, R., Spilotro, G., & Van Westen, C. J. (2016). Rockfall trajectory modeling combined with heuristic analysis for assessing the rockfall hazard along the Maratea SS18 coastal road (Basilicata, Southern Italy). *Landslides*, 13, 985-1003.
- Sigl, R. (1977). *Ebene und Sphaerische Trigonometrie mit Anwendungen auf Kartographie, Geodäsie und Artonomie*, Herbert Wichman Verlag, Karlsruhe, 50-55, 473.
- Stewart I (2009). *Timing The Infinite: The Story of Mathematics*, Quercus, 336s.
- Ulvi, A., Varol, F. i., Yiğit, A. Y. (2019). 3D modeling of cultural heritage: the example of Muiy Mubarek Mosque in Uzbekistan (Hz. Osman's Mushafi). *International Congress on Cultural Heritage and Tourism (ICCHT)*, 115-123, Bishkek, Kyrgyzstan.
- Van Brummelen, G. (2012). *Heavenly mathematics: The forgotten art of spherical trigonometry*. Princeton University Press.
- Yakar, M (2011). Using close range photogrammetry to measure the position of inaccessible geological features. *Experimental Techniques*, 35(1), 54-59.



© Author(s) 2023.

This work is distributed under <https://creativecommons.org/licenses/by-sa/4.0/>